

Certificate of Verification

I, Dr. David Kutteneuler, residing at St. Pauls Platz 1 Munich, Germany, hereby state that I am well acquainted with the German and English languages and that, to the best of my knowledge, the attached document is a true and complete translation of the German priority document DE10210536.7 into the English language.

Munich, June 30, 2010

A handwritten signature in black ink, appearing to read 'D. Kutteneuler', is written over a horizontal line.

Dr. David Kutteneuler

Agent for the preventive therapy after acute stroke

Description

The invention refers to agents for the preventive anti-infective therapy after acute stroke in order to inhibit infections following a stroke event. Infections are severe complications, which commonly occur in the early phase after acute stroke and have a negative prognostic influence. The agents inventively being employed in pharmaceutical preparations are anti-infective agents, in particular antibiotics, which are employed for the prevention and therapy of pneumonias, infections of the urinary tract and sepsis in mammals, in particular in useful animals and domestic animals and also and in particular in the human. The invention moreover refers to agents for the immunomodulating therapy after acute stroke, this approach having the aim to prevent infections after stroke. The agents employed according to the invention in pharmaceutical preparations constitute substances with an immunomodulating potential.

The acute stroke patient, besides the direct consequences of stroke, which may range between transitory or permanent neurological failures and death for reason of encephalothlipsis, in the acute and early remission phase is particularly endangered by infections. Infections, especially pneumonias, constitute the major cause of lethality in stroke (Henon et al. 1995, List of References following the examples). Thus, 21-65% of acute stroke patients develop infections and 10-22% develop pneumonias (Davenport et al. 1996, Castillo et al. 1998, Johnston et al. 1998, Grau et al. 1999, Georgilis et al. 1999, Langhorne et al. 2000). The comparison with the incidence of nosocomial infections occurring in an average of 7-10% of all patients (Bucher 2000) and about 3% of postoperative patients (Smyth & Emmerson 2000), particularly underlines the very high frequency of infections in acute stroke patients. In a systematic investigation, it was shown, that the risk of infection is highest at the first and second day after stroke (Grau et al. 1999).

Moxifloxacin belongs to the class of the fluoroquinolones and comprises the active component *1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-[(4aS,7aS)-octahydro-6H-*

pyrrolo[3,4-b]-pyridin-6-yl]-4-oxoquinoline-3-carboxylic acid, which until now was mainly used against infections of the respiratory tract (Zhanel et al. 2002).

Mezlocillin belongs to the class of the acylaminopenicillins and comprises the active component *(2S,5R,6R)-3,3-dimethyl-6-((R)-2-[3-(methylsulphonyl)-2-oxo-1-imidazolidin-carboxamido]-2-phenylacetamido)-7-oxo-4-thia-1-azabicyclo[3.2.0]heptan-2-carboxylic acid*.

Sulbactam belongs to the class of the β -lactamase inhibitors and comprises the active substance *(2S,5R)-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptan-2-carboxylic acid-4,4-dioxide*.

Mezlocillin in combination with Sulbactam nowadays is predominantly used for the treatment of systemic or local (multiple) infections and as well for the perioperative short-term prophylaxis (Wright 1999).

The invention has the object, to provide medicinal practice with agents, which are suitable for the preventive therapy after acute stroke. This object is achieved by the purposeful application of anti-infective agents. The invention also has the object, to provide medicinal practice with agents, which are suitable to counteract the immunodepression rapidly setting in after acute stroke and being coincided with a consecutive susceptibility to infection.

This object according is achieved by the employment of immunomodulating substances, e.g. of cytokines, inhibitors of the sympathetic nervous system (SNS), inactivated Parapox ovis virus particles or endotoxin binders (rBPI21). According to the invention, medicines for the immunomodulating therapy after acute stroke have been developed, these medicines comprising e.g. cytokines or inhibitors of the SNS, inactivated Parapox ovis virus particles or rBPI21 as active components.

According to the invention, medicines for the preventive therapy after acute stroke have been developed, which - as active components - contain anti-infective substances and/or immunomodulating substances for the prevention and therapy of pneumonias, infections of the urinary tract and sepsis, these active components in particular being selected from the classes of cephalosporines and/or penicillin, tetracyclines, aminoglycosides, lincosamines,

glycopeptides, macrolids, carbapenems, oxazolidinones, streptogramins, linezolides from the fluoroquinolones (here in particular from moxifloxacin), from cytokines (interleukins, interferons), inhibitors of the sympathetic nervous system (beta-blockers, alpha-sympathomimetics), endotoxin binders (lipopolysaccharide binding protein = LBP, BPI, rBPI21) and from inactivated Parapox ovis virus particles in pharmaceutical preparations.

It was moreover found, that a cellular-functional immunodepression, which is mainly mediated by the SNS, occurs after stroke, wherein this immunodepression is such severe, that it may lead to the spontaneous development of severe, prognosis-determining infections. The pharmacological suppression of the SNS, the binding of endotoxins, an immunostimulation or the administration of cytokines (within the first 0 to 7 days) can stop this immunodepression and prevent the development of infections. Moreover, an early-stage preventive therapy with anti-infective agents (within the first 0 to 7 days) can lead to a reduction of lethality and morbidity in acute stroke.

According to the invention, a prolonged and severe immunosuppression with early onset (within the first 12 hours) has been recognized in a commonly accepted animal model of stroke (Mouse; occlusion of the A. cerebri media (MCAO) – Hata et al. 1998). This immunosuppression becomes obvious on the cellular level on the one hand in the form of a lymphopenia of the T-, B- and NK-cells and on the other hand in functional deficits of monocytes and lymphocytes, among other factors caused by a reduced secretion of the pro-inflammatory cytokine IFN- γ . The immunosuppression is mainly mediated by an overactivation of the SNS.

In the course of 2 to 4 days after stroke, a severe infection spontaneously arises in the animals. This infection is a bacterial sepsis and pneumonia with a lethal outcome in about 60% of the animals. By the inventive early administration of a preventive anti-infective therapy (in case of moxifloxacin within the first 24 h), lethality are drastically reduced. In a further aspect of the invention, it is as well possible to prevent the development of infections by a SNS blockade by means of beta-blockers or by the administration of IFN- γ .

The preventive therapy with anti-infective agents (e.g. moxifloxacin) within the first 7 days after the onset of disease constitutes a novel therapeutic approach in the treatment of stroke patients, this approach being suitable to reduce the lethality and morbidity of this disease.

This in particular provides an effective therapeutic approach for the prevention of the commonly occurring complications of infections.

These complications cause a delayed acute-stationary progress and thus prevent the necessary and effective early-stage rehabilitation. These complications are to a large extent responsible for the relatively high lethality after acute stroke (Henon et al. 1995). These complications lead to fever, another independent risk factor for a negative prognosis after acute stroke (Castillo et al. 1998). The treatment of these complications thus reduces lethality and morbidity after acute stroke.

According to the invention, patients suffering from acute stroke are preventively treated within 24 hours after the onset of symptoms with an antibiotic (moxifloxacin) or several anti-infective agents and are thus prevented from infections (like e.g. pneumonias, infections of the urinary tract and sepsis). This reduces direct and indirect lethality and morbidity. The preventive anti-infective therapy is to be continued for 1-7 days.

The basic idea of the invention is to be found in the use of known means for a novel purpose and in a combination of known elements: the anti-infective agents / antibiotics and a novel effect – their employment for affecting complications of infection by means of an early, preventive anti-infective therapy after acute stroke, this concept in its novel combined effect offering an advantage and the desired result, which are to be found therein, that one can now dispose of means for a preventive therapy leading to a reduction of lethality and morbidity after acute stroke.

The invention is moreover directed to the use of anti-infective agents / antibiotics for the preventive anti-infective therapy after acute stroke and for the production of medicines and/or pharmaceutical preparations for the preventive anti-infective therapy after acute stroke.

This comprises their use for the production of medicines for the preventive therapy of pneumonias, infections of the urinary tract and sepsis after acute stroke.

By “early preventive anti-infective therapy after acute stroke” we mean, that the treatment is started within 72 hours after the stroke event.

A further aspect of the invention is given by the immunomodulating therapy with cytokines (e.g. IFN- γ), inactivated parapox viruses (e.g. inactivated Parapox virus ovis), the binding of endotoxin (e.g. by rBPI21) or the blockade of the SNS (e.g. with beta-blockers such as propranolol) within the first 7 days after the onset of the disease. This aspect constitutes a novel therapeutic approach in the treatment of stroke patients, which is suitable to reduce lethality and morbidity in this disease. In particular, this provides an effective therapeutic approach for the prevention of the frequently occurring complications of infections. These complications lead to a delayed acute-stationary progress, thus preventing the necessary and effective early-stage rehabilitation. These complications are major causes for the relatively high lethality after acute stroke (Henon et al. 1995). These complications moreover lead to fever - an independent risk factor for a negative prognosis after acute stroke (Castillo et al. 1998). The prevention of these complications thus reduces lethality and morbidity after acute stroke.

According to the invention, acute stroke patients are treated within 72 h after the onset of symptoms with a cytokine and/or by a pharmacological blockade of the sympathicus and/or by a pharmacological binding of endotoxin or by immunostimulation, thereby preventing the development of a state of immunodepression and offering protection from infections (like e.g. pneumonias, infections of the urinary tract and sepsis). This allows for a reduction of the direct and indirect lethality and morbidity. This immunomodulating treatment is to be performed from day 0 - 7 after the acute stroke event, this being dependent on the clinical study performed in the individual case.

The invention's basis idea is given by the use of known means for a novel purpose and by a combination of known elements - the cytokines, immunostimulating agents and anti-sympathicotonic agents - and a novel effect - their application for the prevention of an immunodepression after acute stroke - which in their novel total effect result in a benefit and in the desired success, which is to be found therein, that one can now dispose of means for a preventive therapy leading to a reduction of lethality and morbidity after acute stroke.

The invention is further directed to the use of cytokines and medicines for blocking the SNS, to endotoxin binders and immunostimulating agents for the immunomodulating therapy after acute stroke and as well to the production of medicines and/or pharmaceutical preparations for the immunomodulating therapy of acute stroke.

This comprises their use for the production of medicines for the mitigation or termination of an immunodepression after acute stroke. By “early immunomodulating therapy after acute stroke” we mean, that the therapy is started within 1 week after the stroke event.

The invention refers in one aspect to the use of an anti-infective agent and/or an immunomodulating agent for the production of a medicine and/or a pharmaceutical preparation for the preventive anti-infective therapy after acute stroke. What is developed according to the invention are medicines and pharmaceutical preparations for the preventive therapy after acute stroke, these medicines/preparations containing as active components anti-infective agents for the prevention and therapy of pneumonias, infections of the urinary tract and sepsis, in particular being selected from the classes of the cephalosporines and/or penicillin, tetracyclines, aminoglycosides, lincosamines, glycopeptides, macrolids, carbapenems, oxazolidinones, streptogramins, linezolides and fluoroquinolones - here in particular Moxifloxacin - as well as cytokines (interleukins, interferons), inhibitors of the sympathetic nervous system (beta-blockers, alpha-sympathomimetics), endotoxin binders (lipopolysaccharide binding protein = LBP, BPI, rBPI21) and inactive Parapox ovis virus particles in pharmaceutical preparations.

The preventive therapy with anti-infective agents (such as moxifloxacin) within the first 7 days after the onset of disease constitutes a novel therapeutic approach in the treatment of stroke patients, which is suitable to reduce lethality and morbidity of this disease.

The frequently occurring complications of infections cause a delayed acute-stationary progress and thus prevent the necessary and effective early-stage rehabilitation. In addition, they are to a large extent responsible for the relatively high lethality after acute stroke (Henon et al. 1995) and moreover give rise to fever, which as an independent risk factor results in a negative prognosis after acute stroke (Castillo et al. 1998). The treatment of these complications thus reduces lethality and morbidity after acute stroke.

According to the invention, patients suffering from acute stroke are treated the earliest possible, at the latest however within 48 hours after the onset of symptoms, with an anti-infective agent, in particular with a broad-spectrum antibiotic, especially with an antibiotic (Moxifloxacin) or a multiplicity of anti-infective agents. This reduces the direct and indirect

lethality and morbidity. The preventive anti-infective therapy should be performed for a period of 1-7 days. The preferred dose is adapted to the anti-infective agent's/agents' respective, common daily dose, which is effective for the treatment of the above mentioned infections. The anti-infective agent is preferably administered on the oral or parenteral route. Moxifloxacin is administered in a dose of 400 mg one time per day on the oral or parenteral route.

The anti-infective medicine to be employed may contain one or several antibiotics in a pharmaceutically preparation. In a specific embodiment of the invention, several different anti-infective agents are combined.

Preferred are the antibiotics from the classes of the cephalosporines and/or penicillin, tetracyclines, aminoglycosides, lincosamines, glycopeptides, macrolids, carbapenems, oxazolidinones, streptogramins, linezolides and/or fluoroquinolones. Of these antibiotics, broad-spectrum antibiotics, especially fluoroquinolones and carbapenems are particularly preferred. Most preferred is the application of moxifloxacin (*1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-[(4aS,7aS)-octahydro-6H-pyrrolo[3,4-b]pyridin-6-yl]-4-oxoquinoline-3-carboxylic acid*), because moxifloxacin is effective in respect to the infective germs emerging in stroke patients and has to be administered just once a day and its safety

The inventive use of an anti-infective agent for the production of medicines and/or pharmaceutical preparations for the preventive anti-infective therapy after acute stroke is useful for mammals, in particular useful animals and domestic animals, but also and in particular for the human. Besides the major use in the field of human medicine, these substances may also be used to produce products and medicines for veterinary medicine.

Preferred is the use of an anti-infective agent in order to provide protection from infections - pneumonias, infections of the urinary tract and/or sepsis - after acute stroke.

According to the present invention, one employs an immunomodulating agent for the production of a pharmaceutical formulation, which immunomodulating agent is characterised in that it may be selected from a group comprising cytokines and/or inhibitors of the SNS, wherein specifically the employment of interferons, interleukins and beta-receptor blockers is preferred. The most preferred substances/agents are the β -blocker propranolol, the interferon

IFN- γ , the endotoxin binder rBPI-21, the lipopolysaccharide binding protein = LBP and inactivated Parapox ovis virus particles.

According to the invention, the immunomodulating agent(s) are used for the production of a medicine or pharmaceutical formulation, which is useful for the preventive anti-infective therapy after acute stroke in mammals, in particular in useful animals and domestic animals, but also and in particular in the human.

The invention also intends to use one or more anti-infective agents and one or more immunomodulating agents for the production of pharmaceutical preparations for the preventive anti-infective therapy after acute stroke, wherein the anti-infective agent can be selected from the group comprising broad-spectrum antibiotics, and wherein the immunomodulating agent can be selected from the group comprising inhibitors of the SNS, cytokines, endotoxin binders and inactivated Parapox ovis virus particles. In a preferred form, the anti-infective agent is moxifloxacin and the immunomodulating agent is IFN- γ .

The invention may also be realized in the form of a kit, which comprises in a separate or combined form a pharmaceutical composition containing an immunomodulating agent, which can be selected from the group of the cytokines and/or the inhibitors of the SNS and an anti-infective agent, which can be selected from the group of the cephalosporines and/or penicillin, tetracyclines, aminoglycosides, lincosamines, glycopeptides, macrolids, carbapenems, oxazolidinones, streptogramins, linezolid and/or fluoroquinolones.

The invention moreover refers to a method for the treatment of patients after acute stroke, this method having the aim to reduce lethality, thereby also being capable to prevent infections after the stroke. In this treatment method, a patient being in need of such a treatment, i.e. a stroke patient, receives an effective dose of the anti-infective agent and/or the immunomodulating agent. Preferably, the respective dose is administered immediately after the stroke event. It is also possible to administer combinations of one or more anti-infective agents and one or more immunomodulating agents. Preferred are the following combinations of one or more broad-spectrum antibiotics and one or more cytokines and/or endotoxin binders and/or Parapox ovis virus particles. During the following days 1-7, one should use the common daily doses of the above mentioned anti-infective agents (1 mg - 100 g) and immunomodulating agents (1 μ g-100 g).

The inventive use of the agents can be realized by known formulations, thus by formulations to date being used for the respective active substances. However, the invention also intends to design novel formulations/routes of application being especially suitable for stroke patients, e.g. intrathecal, nasal, intramuscular, intratracheal, intrabronchial, subcutaneous, intravenous, intraarterial, permucosal, enteral and oral forms.

The inventive use of the agents can be realized in known ways by incorporating them into common formulations like tablets, coated tablets, pills, granules, aerosols, emulsions, suspensions and solutions, this being accomplished under the employment of inert, non-toxic, pharmaceutically acceptable carrier substances or solvents. In this context, the agents' therapeutically effective concentration should correspond to concentrations commonly used for these agents so far; or they should be present in concentrations of about 0,1 to 95 % by weight, preferably of about 0,5 to 90 % by weight of the total composition, i.e. in amounts being sufficient to reach the mentioned dose range. The formulations are e.g. produced by extending the agents with solvents and/or carrier substances, optionally under the employment of emulsifying agents and/or dispersing agents, wherein, e.g. in the case of using water as a diluent, one may optionally use organic solvents as adjuvant substances.

As examples for the adjuvant substances, one may mention e.g. water, non-toxic solvents such as paraffin (e.g. naphtha fractions), plant oils (e.g. peanut / sesame oil), alcohols (e.g. ethanol, glycerol), carrier substances such as natural rock powders (e.g. kaolins, alumina, talcum, chalk), synthetic mineral powders (e.g. highly disperse silica acid, silicates), sugars (e.g. saccharose, lactose and glucose), emulsifying agents (e.g. polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers), dispersing agents (e.g. lignin, sulphite lyes, methylcellulose, starch and polyvinyl pyrrolidone) and lubricants (e.g. magnesium stearate, talcum, stearic acid and sodium sulphate). The application is usually performed preferably via the oral or parenteral route, in particular perlingually or intravenously. In case of the oral application of the medicines according to the inventions, tablets - besides the mentioned carrier substances - may obviously also contain additives like sodium citrate, calcium carbonate or dicalcium phosphate together with different excipients and filling materials like starch, preferably potato starch, gelatine and the like. Moreover one may also use lubricants like magnesium stearate, sodium lauryl sulphate and talcum for forming the tablets. In case of aqueous suspensions, the active substances - besides the above mentioned adjuvants - may

also be mixed with different flavour-improving substances or colorants. In case of the parenteral applications one can employ solutions of the active substances by using suitable liquid carrier materials.

The invention further comprises a kit comprising in a separate form a pharmaceutical composition containing an immunomodulating substance, which may be selected from the above described group of immunomodulating substances and an anti-infective agent, which may be selected from the above described group of the anti-infective agents.

The invention moreover comprises the use of immunomodulating agents and anti-infective agents for the production of a pharmaceutical composition or pharmaceutical medicine containing both substances. Likewise, several substances of both therapeutic groups may be present. Preferred are broad-spectrum antibiotics as well as inhibitors of the SNS and/or cytokines and/or endotoxin binders and/or Parapox ovis virus particles.

More preferred combinations are fluoroquinolones and carbapenems and/or cytokines. Most preferred are moxifloxacin and IFN- γ .

The invention will now be further illustrated by means of examples with reference to the figures, however without being limited to them. The figures show:

Figure 1: Bacteraemia and pneumonia 3 days after experimentally induced stroke.

Figure 1B: Cell separation by FACS-analysis from blood (a-c), spleen (d-f) and thymus (g), and tests of lymphocyte functions for monocytes (h) and T-cells (i).

Figure 2A: Hyper- and hypothermia after experimental stroke and its effective prevention by an early preventive therapy with an anti-infective agent.

Figure 2B: Bacteraemia and pneumonia 3 days after experimentally induced stroke.

Figure 3A: Lethality after experimental stroke and its effective prevention by an early preventive therapy with an anti-infective agent.

Figure 3B: Prevention of the lymphopenia (a), the disturbance of lymphocyte function (b) and the development of severe infections (c) by means of a pharmacological sympathetic blockade.

Figure 4: Attenuation of the severity of infection in the lung and in the blood (number of CFU) by the administration of a cytokine (IFN- γ).

Examples

A) Anti-Infective Agent

Example 1A: Mouse model of cerebral ischemia

According to the mouse model of cerebral ischemia mice were operated according to the MCAO. In this operation, the A. cerebri media is occluded for about 60 minutes, leading to a cerebral infarction typical for this model. As a control, we use the so-called Sham-operation, in which the animals are also MCAO-operated, but in which the A. cerebri media is only occluded for about 1 minute. For the rest, the two paradigms are identical. Thus, the perioperative stress can be doubtlessly excluded as a systemic mistake for the following results.

Figure 1A shows that the mice, 3 days after stroke, suffer from a bacteraemia and a pneumonia with a load of bacterial infection (more than 95% *Escherichia coli*) of 2×10^4 or, respectively, of 4×10^6 colony forming units/ml (CFU/ml). In contrast, no infection is found in the control animals (Sham-operated).

The stroke animals develop fever within the first 12 hours (h) after the stroke event and start to exhibit a hypothermic body temperature after further 12 h (Figure 2, vector group). In the further process, the bacteraemia/pneumonia proceed to the clinical picture of a sepsis, which leads to death after 4-6 days (Figure 3, vector group).

Example 2A: Application of an anti-infective agent

By means of a preventive therapy based on the anti-infective agent mezlocillin plus sulbactam, a very early - 0 to 12 h after acute stroke - treatment scheme allowed to prevent both the infections (data not shown) and to significantly reduce the fever/hyperthermia and in particular the lethality after stroke.

Example 3A: Application of Mezlocillin plus Sulbactam

By means of a preventive therapy following the principle of example 2, the employment of mezlocillin and sulbactam in a later - 12 to 24 h after acute stroke - treatment scheme, both the infections were prevented (data not shown) and the fever/hyperthermia and in particular the lethality after stroke were significantly reduced.

Example 4A: Application of Moxifloxacin

By means of a preventive therapy with moxifloxacin in a very early (0-12 h after acute stroke) and as well in a somewhat later (12-24 h after acute stroke) treatment scheme, both the infections (pneumonia, sepsis) were prevented and the fever/hyperthermia and in particular lethality after stroke were significantly reduced (figures 2 and 3).

B) Immunomodulation

Example 1B: Mouse model of cerebral ischemia

According to the mouse model of cerebral ischemia, mice were operated according to the MCAO. In this operation, the A. cerebri media is occluded for about 60 minutes, leading to a cerebral infarction typical for this model. As a control, we use the so-called Sham-operation, in which the animals are also MCAO-operated, but in which the A. cerebri media is only occluded for about 1 minute. For the rest, the two paradigms are identical. Thus, the perioperative stress can be doubtlessly excluded as a systemic mistake for the following results.

Figure 1B shows, that the mice within a half day after stroke exhibit a significant reduction in the number of lymphocytes (more detailed: B-cells, T-cells and NK-cells) circulating in the blood (Fig. 1:a-c). This severe lymphopenia persists at least for 14 days and also affects the spleen and the thymus (Fig. 1:d-g), which constitute essential organs of immunological maturation. Besides the mere numeric decrease, there also occurs a functional impairment with a disturbance of the cytokine secretion capacity after stimulation (Fig. 1: h and j). Besides the disturbance of monocyte function, which manifests as a disturbance of TNF- α secretion for 2 days after LPS-stimulation, one can also detect a lymphocyte alteration after ConA-stimulation; this alteration persisting for 14 days after the stroke and mainly causing a Th1- to Th2-shift (alteration of the IFN- γ / IL4 ratio), which means a loss of pro-inflammatory potential. Also the Sham-operated animals exhibit alterations in the above mentioned

parameters, but these alterations are smaller. B- and NK-cells only slightly decrease in the blood after the OP-stress. In the spleen and in the thymus, no major stress-induced alterations are detected and the functional parameters of the monocytes and lymphocytes remain at the control level. Figure 2 shows that the mice - 3 days after the stroke event - exhibit a bacteraemia and a pneumonia with a load of bacterial infection (more than 95% *Escherichia coli*) of 2×10^5 or, respectively, of 4×10^7 colony forming units/ml (CFU/ml). In contrast to this, no infection is found in the control animals (Sham-operated).

Example 2B: By means of a blockade of the SNS by the administration of propranolol (an unspecific beta-blocker) after experimental stroke in mice, it was possible, to prevent the decrease of T- and NK-cells and to significantly reduce the degree of B-cell lymphopenia (Fig. 2A). Furthermore, the blockade of the SNS by propranolol also prevents the disturbance of function of the remaining monocytes (TNF- α secretion) and of the lymphocytes (IFN- γ / IL-4 ratio) (Fig. 2B). The SNS blockade can also prevent the development of infections in the lung (pneumonia) and in the blood (sepsis) (Fig. 2C).

Example 3B: Application of cytokines

By the application of IFN- γ , it was possible to reduce both the germ number in the lung and - even more pronounced - the germ number in the blood.

References

- Bucher A (2000). Hand hygiene-is hand disinfection the best solution. *Tidsskr Nor Laegeforen* 120:472-5
- Castillo J, Dávalos A, Marrugat J, Noya M (1998). Timing for fever-related brain damage in acute ischemic stroke. *Stroke* 29:2455-2460.
- Davenport RJ, Dennis MS, Wellwood I, Warlow CP (1996). Complications after acute stroke. *Stroke* 27:415-20
- Georgilis K, Plomaritoglou A, Dafni U, Bassiakos Y, Vemmos K (1999). Aetiology of fever in patients with acute stroke. *J Intern Med* 246:203-9
- Grau AJ, Buggle F, Schnitzler P, Spiel M, Lichy C, Hacke W (1999). Fever and infection early after ischemic stroke. *J Neurol Sci* 171:115-20

- Hata R, Mies G, Wiessner C, Fritze K, Hesselbarth D, Brinker G, Hossmann KA (1998). A reproducible model of middle cerebral artery occlusion in mice: hemodynamic, biochemical, and magnetic resonance imaging. *J Cereb Blood Flow Metab* 18:367-75
- Henon H, Godefroy O, Leys D, Mounier-Vehier F, Lucas C, Rondepierre P, Duhamel A, Pruvo JP (1995). Early predictors of death and disability after acute cerebral ischemic event. *Stroke* 26:392-8
- Johnston KC, Li JY, Lyden PD, Hanson SK, Feasby TE, Adams RJ, Faught RE Jr, Haley EC Jr (1998). Medical and neurological complications of ischemic stroke: experience from the RANTTAS trial. RANTTAS Investigators. *Stroke* 29:447-53
- Langhorne P, Stott DJ, Robertson L, MacDonald J, Jones L, McAlpine C, Dick F, Taylor GS, Murray G (2000). Medical complications after stroke: a multicenter study. *Stroke* 31:1223-9
- Smyth ET, Emmerson AM (2000). Surgical site infection surveillance. *J Hosp Infect* 45:173-84
- Wright AJ (1999). The penicillins. *Mayo Clin Proc* 74:290-307
- Zhanell GG, Ennis K, Vercaigne L, Walkty A, Gin AS, Embil J, Smith H, Hoban DJ (2002). A critical review of the fluoroquinolones: focus on respiratory infections. *Drugs* 62:13-59

Claims

1. Use of one or more anti-infective agents and/or one or more immunomodulating agents for the production of a medicine and/or a pharmaceutical preparation for the preventive anti-infective therapy after acute stroke.
2. Use of an anti-infective agent according to claim 1, characterised in that the anti-infective agent contains one or more antibiotics in a pharmaceutical preparation.
3. Use of an anti-infective agent according to claim 2, wherein the antibiotic(s) can be selected from the classes of the cephalosporines and/or penicillin, tetracyclines, aminoglycosides, lincosamines, glycopeptides, macrolids, carbapenems, oxazolidinones, streptogramins, linezolid and/or fluoroquinolones.
4. Use of an anti-infective agent according to one of the claims 1 to 3, characterised in that moxifloxacin (*1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-[(4aS,7aS)-octahydro-6H-pyrrolo[3,4-b]-pyridin-6-yl]-4-oxoquinoline-3-carboxylic acid*) is contained therein.
5. Use of an anti-infective agent according to claim 3 for the production of a pharmaceutical preparation, wherein mezlocillin and sulbactam are used in combination.
6. Use of an anti-infective agent according to one of the claims 1 to 5 for the production of medicines and/or pharmaceutical preparations for the preventive anti-infective therapy after acute stroke in mammals, in particular in useful animals and domestic animals, and in particular in the human.
7. Use of an anti-infective agent according to one of the claims 1 to 6 for the protection from infections, such as pneumonias, infections of the urinary tract and/or sepsis, after acute stroke.
8. Use of an immunomodulating agent according to claim 1, characterised in that the immunomodulating agent is selected from the group comprising cytokines and/or inhibitors of the SNS or endotoxin binders or Parapox ovis virus particles.


9. Use according to claim 8, wherein the immunomodulating agent(s) are selected from the group comprising interferons and beta-receptor blockers.
10. Use according to claim 8, wherein propranolol and/or IFN- γ is/are employed.
11. Use according to one of the claims 7 to 9 for the production of medicines and/or pharmaceutical preparations for the preventive anti-infective therapy after acute stroke in mammals, in particular in useful animals and domestic animals, and in particular in the human.
12. Use of one or more anti-infective agents and one or more immunomodulating agents for the production of a pharmaceutical preparation for the preventive anti-infective therapy after acute stroke according to one of the foregoing claims, wherein the anti-infective agent can be selected from the group comprising cephalosporines and/or penicillin, tetracyclines, aminoglycosides, lincosamines, glycopeptides, macrolids, carbapenems, oxazolidinones, streptogramins, linezolid and fluoroquinolones, and wherein the immunomodulating agent can be selected from the group comprising cytokines (interleukins, interferons), inhibitors of the sympathetic nervous system (beta-blockers, alpha-sympathomimetics), endotoxin binders and inactivated Parapox ovis virus particles. Use according to claim 11, wherein the anti-infective agent is moxifloxacin and the immunomodulating agent is IFN- γ .
13. Kit, comprising in a separate or combined form a pharmaceutical composition containing an immunomodulating agent, which can be selected from the group of the cytokines and/or inhibitors of the SNS and an anti-infective agent, which can be selected from the group of the cephalosporines and/or penicillin, tetracyclines, aminoglycosides, lincosamines, glycopeptides, macrolids, carbapenems, oxazolidinones, streptogramins, linezolid and/or fluoroquinolones.

Abstract

The invention refers to agents and their uses for the preventive therapy after acute stroke, in particular having the aim to prevent infections after stroke. The agents inventively employed in pharmaceutical preparations are anti-infective agents and/or immunomodulating agents, e.g. cytokines and/or inhibitors of the SNS.

Figure 1A:

1 / 7



Legend to Figure 1A:

Lunge - lung

Blut - blood

Kontrolle - control

Pneumonie - pneumonia

Figure 1B:





Figure 2A:

3 / 7



Legend to Figure 2A:

Temperatur - temperature

Vektor - vector

Figure 2B:

4 / 7





Figure 3A:

5/7



Legend to figure 3A:

Vektor - vector

Überlebensfunktionen - survival functions

Kumulatives Überleben - cumulative survival

Mäuse je Gruppe - mice per group

Tage - days

Figure 3B:

6/7




Figure 4:

7/7

